

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 November 2004 (04.11.2004)

PCT

(10) International Publication Number
WO 2004/094474 A1

(51) International Patent Classification⁷: **C07K 16/12**,
A61K 39/40, C12Q 1/68, G01N 33/563

(21) International Application Number:
PCT/GB2004/001619

(22) International Filing Date: 14 April 2004 (14.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0309126.1 17 April 2003 (17.04.2003) GB

(71) Applicant (for all designated States except US): **NEUTEC PHARMA PLC** [GB/GB]; Clinical Sciences Building, Central Manchester Healthcare Trust, Oxford Road, Manchester Lancashire M13 9WL (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BURNIE, James, Peter** [GB/GB]; 1 Greystoke Drive, Alderley Edge Cheshire SK9 7PY (GB). **MATTHEWS, Ruth, Christine** [GB/GB]; 1 Greystoke Drive, Alderley Edge Cheshire SK9 7PY (GB).

(74) Agents: **MCCALLUM, Graeme, David** et al.; Lloyd Wise, McNeight & Lawrence, Highbank House, Exchange Street, Stockport Cheshire SK3 0ET (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CLOSTRIDIUM DIFFICILE FOCUSED ANTIBODIES

(57) Abstract: The present invention is concerned with antibodies specific for and which confer immunity against infection by C. difficile or against a vaccine, methods of identifying same, methods of manufacture of medicaments, and methods of treatment of patients using same. Also provided is a method for determining the efficacy of a vaccine, together with methods of vaccinating a patient, diagnostic test methods and diagnostic test kits.



WO 2004/094474 A1

- 1 -

Clostridium difficile focussed antibodies

The present invention is concerned with antibodies either specific for and which confer immunity against infection by *Clostridium difficile* or antibodies produced after challenge with a *C. difficile* based vaccine, methods of identifying same, and uses of same.

C. difficile is a gram-positive anaerobic bacterium, and is deemed a significant human pathogen causing a spectrum of diseases ranging from mild diarrhoea to fulminant pseudomembranous colitis (PMC) - collectively referred to as *C. difficile* antibiotic-associated diarrhoea (CDAD). CDAD is a common, iatrogenic, nosocomial disease associated with substantial morbidity and mortality, especially in the elderly. Two factors have been assigned main roles in the pathogenesis of CDAD - the suppression of the resident intestinal flora by the administration of antibiotics, and the production by the bacterium of two high molecular weight toxins, exotoxin A and exotoxin B.

The bacterium is endemic in hospitals, and studies have shown that approximately one third of patients receiving antibiotic treatment in acute-care medical wards were colonised by *C. difficile* while in hospital (Kyne, L., *et al.*, 2002, Clin. Infect. Dis. 34(3), pp346-53, PMID: 11774082). Of these patients, over half went on to develop CDAD while the remainder were symptomless carriers. CDAD is a major factor in extension of patient hospital stay times, and estimates suggest that the cost of this disease in the US exceeds \$1.1 billion per year (Kyne, L., *et al.*, *Supra*). Patients suffering from CDAD respond well to a treatment which includes a discontinuation of the inciting antibiotic and treatment with either of the antibiotics metronidazole and vancomycin. However, the use of e.g. vancomycin is one of last resort since it is associated with several problems. Not only may it cause nephrotoxicity, ototoxicity, bone marrow toxicity and the red man syndrome, but the problem with this treatment regime is that the CDAD often returns after successful treatment of the initial episode, and this reoccurrence represents a serious clinical problem.

- 2 -

Additionally, there is evidence that *C. difficile* is becoming resistant to metronidazole and partially resistant to vancomycin, demonstrating the need for new alternatives in the treatment of CDAD.

5 Exotoxins A and B which are produced by pathogenic strains of the bacterium are cytotoxic, enterotoxic and proinflammatory, and are considered to be the main virulence factors of this non-invasive microorganism. However, not all infections with toxigenic strains result in disease, prompting the search for additional virulence factors. Bacterial surface expressed antigens represent candidate virulence factors, and are also considered
10 important since such proteins likely mediate the essential functions such as adhesion to the epithelial layer of the gut in the first step of colonization or interaction with mediators of local immunity. In common with many other bacteria, *C. difficile* expresses a crystalline or paracrystalline surface layer (S-layer) on the outer cell surface. Such S-layers comprise proteins or glycoproteins forming a regularly arranged lattice on the external surface of the
15 bacterium, and have previously been shown to be essential for the virulence of pathogens such as *Aeromonas salmonicida* and *Campylobacter fetus*. In contrast to most bacteria which comprise one S-layer, *C. difficile* is known to comprise two superimposed paracrystalline S-layers, each composed of a glycoprotein subunit which varies slightly in apparent molecular weight among different *C. difficile* strains. Most strains of *C. difficile*
20 express two major S-layer proteins (SLPs), one of 32-38 kDa (low-MW SLP) and a second of 42-48 kDa (high-MW SLP). The low-MW SLP appears to be immunodominant and is the antigen most commonly recognised by patients suffering from CDAD, and is the only antigen recognised in EDTA extracts of bacteria by antisera raised in rabbits against whole *C. difficile* cells (Calabi, E. et al., 2001, Mol. Microbiol., 40(5) p1187-99, PMID:
25 11401722).

During the course of microbial infection various adaptive strategies are employed by the immune system. One such strategy, and arguably the most important, is the production of

- 3 -

an antibody response. Antibodies capable of binding antigens displayed by the infectious agent are produced and bind to and allow killing of the microorganism through complement activation, recruitment of macrophage and through direct interaction with the microbe itself. The therapeutic efficacy of antibodies capable of binding a given antigen varies and this is reflected in the fact that antibody production by the immune system matures during the course of an infection and becomes focussed in the case of a patient successfully fighting off an infection.

The antibody response is elicited by the B cell repertoire where individual B cells each produce structurally diverse antibody molecules. The actual size of this B cell/antibody repertoire is unknown, but it is estimated that the random clonal frequency of reactivity for a given antigen may be as high as 1 in 100,000 in cultured B cells (Nobrega, A., et al., Eur J Immunol. 1998 Apr;28(4):1204-15; PMID: 9565360). During the course of infection, antibodies capable of binding the pathogen are selected for by changes in the B cell population resulting in key antibodies being produced in large numbers. The mechanisms for these changes include clonal expansion, isotype switching, and somatic mutation of immunoglobulin variable regions. B cells responsible for generating antibodies which are able to bind a pathogen multiply, thus skewing the B cell repertoire and changing the proportions of B cells.

Non-antibiotic based therapeutic regimes for the treatment/prevention of *C. difficile* infection are based upon vaccination and passive immunization. Vaccination treatment comprises administering to a patient either a nucleic acid sequence encoding an immunogenic fragment of the *C. difficile* surface layer protein or a variant or homologue thereof, or an equivalent polypeptide fragment (as disclosed in WO 02/062379). Passive immunotherapy is typically achieved by administering to a patient a monoclonal antibody specific to an immunogen produced by a pathogen. In general, passive immunotherapy is particularly effective in treating immunocompromised patients who are unable to respond

- 4 -

to vaccination, and to patients who need immediate therapy and cannot wait for vaccination to take effect. In the case of a *C. difficile* infection, passive immunization relies on the administration to a patient of toxin-neutralizing polyclonal immune globulin, (as disclosed in WO 99/2030.4), or antibodies raised against the whole bacterium and the
5 toxins (as disclosed in WO 96/07430).

Thus, as can be seen from the prior art, in order to effect treatment of the patient, an immunogen is first isolated and purified, administered to test animals, and cells producing antibody specific against the immunogen cloned. The range of antibodies produced by the
10 clones can then be tested for their therapeutic efficacy and a single monoclonal antibody selected which is then administered to patients to effect passive immunotherapy.

There are, however, several problems associated with current passive immunotherapy regimes aimed at treating *C. difficile* infections. For example, passive immunotherapy
15 requires that there are survivors of the *C. difficile* infection and patients who have been vaccinated. Each batch of antibody can be different leading to difficulties associated with standardisation and administration of the immunotherapeutic reagent. In addition, the problem of inadvertent administration to a patient of adventitious agents (e.g HIV, HBV, HCV, or as yet unidentified agents) is a real one, and up-to-date screening of any
20 immunotherapeutic reagent is required. Finally, the strain variability exhibited by *C. difficile* means that a given antibody may only be useful against certain strains of the bacterium and not against others.

Obviously the techniques involved are somewhat complex, inconvenient, expensive and
25 time consuming. In general they require that an immunogen is isolated from the infecting pathogen and used to generate antibody. Simply isolating the immunogen can be extremely difficult and time consuming, particularly if it comprises carbohydrate or complex non-linear epitopes (i.e. epitopes having secondary, tertiary and/or quaternary structural

- 5 -

features) which cannot be synthesised *in vitro*, and are impossible to isolate and produce as an antigen for use as e.g. a vaccine. The SLPs of *C. difficile* contain a glycoprotein subunit which varies in molecular weight between species. It may be the case that *C. difficile* epitopes are only produced *in vivo* and are not synthesised *in vitro*, consistent with
5 for example *Neisseria gonorrhoeae* infections (the causal agent of gonorrhea) where specific antigens are only expressed upon infection of a host. Such antigens fall into the general class of cryptoantigens. Furthermore, *C. difficile* may display highly labile antigens which are difficult to work with since during use they simply degrade and the epitope they display is lost. With regard to this range of epitopes/antigens whose identification and/or
10 *in vitro* use is of great difficulty or impossible, the present invention overcomes these disadvantages and provides a solution by providing antibodies whose CDR regions have been generated in response to *C. difficile* epitopes during antibody responses of patients infected with *C. difficile*.

15 In addition, the prior art typically has to attempt to achieve an equivalent of affinity maturation of antibodies by first synthesising a set of candidate antibodies specific to an antigen, testing them for their binding characteristics and then modifying the sequences of the candidates in order to optimise the binding. A thorough attempt at affinity maturation (i.e. optimising antibody binding) can require the synthesis of thousands of
20 different antibodies, which can be costly and time-consuming. Because the antibodies of the present invention are obtained from patients who have either been infected by a pathogen displaying the antigen or who have been vaccinated with an antigen, they have by their very nature and definition already undergone affinity maturation, as is most clearly demonstrated by the sequences of their CDR3 regions of the variable heavy and variable
25 light chains (i.e. the CDR-H3 and CDR-L3 regions).

- 6 -

According to the present invention there is provided an antibody or an antigen binding fragment thereof having the CDR-H3 sequence selected from the group consisting of: SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, and SEQ ID NO: 31.

- 5 Also provided according to the present invention is an antibody or an antigen binding fragment thereof having the CDR-L3 sequence selected from the group consisting of: SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO: 34.

- 10 Also provided according to the present invention is an antibody or an antigen binding fragment thereof having a CDR-H3 sequence selected from the group consisting of: SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, and SEQ ID NO: 31, and a CDR-L3 sequence selected from the group consisting of: SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO: 34.

- 15 According to the present invention there is also provided a method for identifying candidate sequences of at least the CDR3 region of antibodies specific against at least one antigen produced by *C. difficile* during an infection or against a vaccine, comprising the steps of:

- 20 (i) with B cells isolated from at least one patient who has been infected by *Clostridium difficile* or administered said vaccine, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells; and
- (ii) correlating said sequenced at least the CDR3 regions of the VH and/or VL coding regions of said B cells from said at least one patient to
- 25 identify a set of candidate sequences for at least a CDR3 region of antibodies specific against said at least one antigen produced by *Clostridium difficile* or against said vaccine, each of said set of candidate CDR3 sequences or a sequence having at least 80%

- 7 -

homology therewith occurring in total at a frequency of at least 1 percent in the set of sequences determined at step (i).

5 Examples of patients used as a source of B cells in such a method are humans and other mammals such as mice, rabbits, rats, baboons, monkeys and apes.

In certain embodiments of the invention, step (i) may comprise the steps of:

- (i)(a) isolating B cells from at least one patient who has been infected by *Clostridium difficile* or administered said vaccine; and
- 10 (i)(b) sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells;

The sequences occurring in total at a frequency of at least 1 percent in the set of sequences determined at step (i) may be the candidate CDR3 sequences or a sequence having at least
15 80% homology, for example 85, 90, 95, 96, 97, 98 or 99% homology therewith. Sequence homology is as determined using the BLAST2 program (Tatusova TA *et al.*, FEMS Microbiol Lett. 1999 May 15;174(2):247-50; PMID: 10339815) at the National Center for Biotechnology Information, USA (www.ncbi.nlm.nih.gov) with default parameters.

20 For example, the sequences occurring in total at a frequency of at least 1 percent in the set of sequences determined at step (i) may be the candidate CDR3 sequences or a sequence having 1 or 2 amino acid changes therefrom. In general, the frequency of a dominant sequence is added together with the frequency of any sequence showing >80% homology, such that any sequence exhibiting a frequency of 1 % or greater is then deemed a candidate
25 sequence. For example, a first sequence may occur at a frequency of 0.7 percent, and first, second, third and fourth sequences each having a single amino acid change there from each occur at a frequency of 0.1% - the total occurrence is therefore 1.1% and the dominant

- 8 -

antibody sequence (occurring at a frequency of 0.7%) is therefore a candidate CDR3 sequence.

5 This is fundamentally different to prior art methods of making a library of antibodies in a vector such as a phage library where antibodies are isolated by panning and which require binding with a specific (known) antigen in a structurally- and charge-specific state. An analogy by which the two can be compared would be to say that the prior art making of a library and subsequent panning is like going into a library, picking up a book and hoping that it is the right one. In comparison, the present invention is like going into a library,
10 reading all of the books in the library and finding that one book is dominant - that there are many copies of it - and that it is relevant to the disease to be treated.

In addition, the prior art use of library systems has a number of significant problems - in particular, some antibodies produced by a library may cause the death of the organism
15 expressing them and therefore they simply cannot be detected. This is not a particular problem when e.g. looking for antibodies specific to cancers, but when one is searching for antibodies specific to an antigen from a pathogen which might be homologous to one produced by the host expression system (e.g. *Escherichia coli*) then important antibodies cannot be expressed. The use of e.g. *E. coli* to express libraries of e.g. human antibodies
20 also suffers from the problem of codon usage - codons used by humans for specific amino acids can frequently not be the optimum ones for the same amino acid in *E. coli* or other host systems. This means that an important antibody might not be expressed (or at least not in sufficient quantities) since the codons in its sequence are highly inefficient in *E. coli*, resulting in the *E. coli* being unable to read through and express it in full. Codon
25 optimisation of antibody libraries is obviously not an option since the libraries would first have to be sequenced, which defeats the main advantages of using libraries. Since the present invention sequences antibodies directly from a patient, it avoids this problem.

- 9 -

The correlation step (ii) may also correlate the occurrence of the candidate sequences against their occurrence in patients who have not been infected with *Clostridium difficile* or administered the vaccine, sequences only being determined to be candidate sequences if they or a sequence having 80% homology to them occur in total at a frequency of less than 1 percent in the non-infected/vaccinated patients.

The at least one antigen produced by the micro-organism may of course be an immunogen.

The present invention provides unique opportunities for understanding antibody responses to infection which were not previously available, and provides novel diagnostic and therapeutic opportunities.

In particular, since the methods of the present invention bypass the need to isolate an antigen from *C. difficile*, the difficulties associated with identifying e.g. complex non-linear epitopes, epitopes containing carbohydrate, or cryptoantigens are avoided. Instead the methods facilitate the determination of the identity of antibodies specific against one or more antigens produced by the bacterium. As is explained below, the methods of the present invention allow the identification of therapeutically effective antibodies and can allow the identification of the most important parts of antibody sequences. This is particularly evident when the B cells of a given patient are sampled at different time points during the course of an infection by *C. difficile* - as the patient's immune system becomes focussed on producing therapeutically effective antibodies against the bacterium, a focussing of antibody sequences is observed, and the development of variable and non-variable regions within the CDR parts of the VH and/or VL sequences is observed. In addition, the methods of the present invention can allow the identification of antibodies best suited to the treatment of *C. difficile* infection in particular groups of patients, such as age, sex and racial groups. Similarly by comparing the antibody sequences of patients who have recovered from infection with *C. difficile* with those that have not recovered

- 10 -

from infection, ineffective antibody sequences (which might be produced by both sets of patients) can be identified.

5 The B cells isolated from the at least one patient can be peripheral B-cell lymphocytes (PBLs), and can be isolated from a blood sample from the at least one patient. B-cells can also be isolated from other sources and the present invention extends to their use. For example, B-cells can be isolated from spleen.

10 Antibody CDR sequences are determined as detailed in the "Experiments" section using standard techniques and definitions of their start and end.

As is detailed below, the antibody sequences are determined from patients with *C. difficile* infections. These sequences can be derived from B cell immunoglobulin mRNA and hence reflect expressed antibodies and repertoires therein. The sequences determined according
15 to the present invention need not be the sequences of whole antibody molecules – they comprise at least the VH and/or VL sequences which specify the regions responsible for antigen binding.

From the nucleotide sequences determined by the initial sequencing, putative amino acid
20 sequences for the VH and/or VL regions can be determined using standard algorithms and software packages (e.g. see www.mrc-lmb.cam.ac.uk/pubseq/, the Staden package and Gap4 programs; Rodger Staden, David P. Judge and James K. Bonfield. Managing Sequencing Projects in the GAP4 Environment. Introduction to Bioinformatics. A Theoretical and Practical Approach. Eds. Stephen A. Krawetz and David D. Womble.
25 Human Press Inc., Totawa, NJ 07512 (2003); Rodger Staden, David P. Judge and James K. Bonfield. Analysing Sequences Using the Staden Package and EMBOSS. Introduction to Bioinformatics. A Theoretical and Practical Approach. Eds. Stephen A. Krawetz and David D. Womble. Human Press Inc., Totawa, NJ 07512 (2003)). These can be further

- 11 -

characterised to determine the CDR (Complementarity Determining Region) parts of the VH and/or VL sequences, particularly CDR1, CDR2 and CDR3. Methods for determining the putative amino acid sequences and identifying CDR regions are well known and detailed below.

5

As well as the VH and/or VL amino acid sequences, several pieces of additional information can be used in the correlation step:

10

- (i) the strain of *C. difficile* causing the patient's infection;
- (ii) the time point during the infection process at which antibody sequences were sampled
- (iii) patient details - none or more of: sex, race, and age; and
- (iv) the range of complementarity determining regions (CDR) ie the variable regions of the antibody that undergo direct antigen binding/contact.

15

Thus in a method according to the present invention, B cells can be used which have been isolated from the at least one patient at a plurality of time points during infection of the at least one patient by *C. difficile* (or post-vaccination), correlation step (ii) correlating the time point during infection of the at least one patient by *C. difficile* at which the B cells are isolated.

20

25

Similarly, in a method according to the present invention, B cells can be used which have been isolated from at least two patients, at least one of whom has recovered from infection by *C. difficile*, and at least one of whom has not recovered from infection by *C. difficile*, correlation step (ii) correlating the recovery of the at least two patients from infection by *C. difficile*.

- 12 -

Similarly, in a method according to the present invention, B cells can be used which have been isolated from at least two patients, the patients being infected by strains of *C. difficile* producing the at least one antigen, correlation step (ii) correlating the sequenced at least the VH and/or VL coding regions of the B cells to identify a set of candidate sequences for
5 antibodies, each of which is specific against at least one shared antigen produced by the different strains of *C. difficile* or is specific against different antigens produced by the different strains of *C. difficile*.

The sequencing of the VH and/or VL regions can also be used to identify the surrounding
10 antibody framework, and this can be used to determine the antibody isotype. This can then be used in the correlation step to determine whether specific antibody isotypes are particularly useful.

By the term "therapy" is meant any treatment which is designed to cure, alleviate, remove
15 or lessen the symptoms of, or prevent or reduce the possibility of contracting any disorder or malfunction of the human or animal body

The invention is suitable for identifying antibody sequence useful in treating *C. difficile* infection.

20 Using the abovementioned additional information fields in the correlation step (ii), it is easy to perform the correlation using e.g. only CDR sequences, CDRs from either a given patient or a given infection or both, CDRs from a given site or patient group, or CDRs from different sampling times over the course of infection.

25 These correlations are CDR-specific because these are the hypervariable regions responsible for antigen binding and sequence diversity of the antibody repertoire (although

- 13 -

the framework regions can be used in the correlation step as well to reveal the antibody isotype).

The present invention is useful in a wide range of applications, e.g. antibody therapy, and vaccine studies.

Antibody Therapy:

This involves the passive transfer of antibody to non-immune individuals (e.g. patients undergoing chemo-/radio-therapy, immunosuppression for organ transplantation, immunocompromised due to underlying conditions such as diabetes, trauma etc, also the very young or very old).

The present invention can be used to determine the sequences of antibodies conferring immunity by looking for over-represented VH and/or VL sequences in patients who have overcome infection. These protective antibodies can be re-synthesised at the genetic level, over-expressed in *E. coli* (or other expression systems) and purified. The resultant purified recombinant antibody can then be administered to patients as a passive immunotherapy. Antibodies can also be ordered from commercial suppliers such as Operon Technologies Inc., USA (www.operon.com) by simply supplying them with the sequence of the antibody to be manufactured.

Therapeutically useful sequences can be identified in a number of ways which can be used in the correlation step (ii):

- 1) By looking for antibody sequence over-representation in patients who have recovered from *C. difficile* infection - B cells that produce antibodies capable of pathogen binding undergo clonal expansion,

- 14 -

hence high frequency antibody sequences are most likely to bind pathogen and confer immunity.

- 2) By following alterations in the antibody repertoire over the course of *C. difficile* infection. During infection, antibodies undergo a maturation process to improve pathogen binding and this is characterised by sequence alterations. Also, B cell clonal expansion is more prevalent in the final stages of infection where the infection is cleared. The most frequent antibodies in the repertoire are chosen as candidates for immunotherapy. Following the maturation process the candidate antibody will demonstrate which key amino acid residue alterations improve antigen binding and this information can be used to improve antibody design.
- 3) Analysis of the repertoires from different patients with *C. difficile* infection can identify shared and identical protective antibodies. These antibodies are attractive choices for immunotherapy as their occurrence in different patients suggests a strong positive selection in their favour; hence an important role in antibody-based immunity.
- 4) Analysis of the repertoires from patients infected with different strains of *C. difficile*. In this situation, if a common antibody sequence is found in the repertoires for both strains, the antibody may be useful in treating both strains of the bacterium, ie displaying a broad spectrum.
- 5) Analysis of the repertoires for affinity maturation of sequences.

- 15 -

Vaccination Studies:

Vaccination protects against infection by priming the immune system with pathogen-derived antigen(s). Vaccination is effected by a single or repeated exposures to the pathogen-derived antigen(s) and allows antibody maturation and B cell clonal expansion without the deleterious effects of the full-blown infectious process. T cell involvement is also of great importance in effecting vaccination of patients. The present invention can be used to monitor the immunisation process with experimental *C. difficile* vaccines. Subjects are given the experimental vaccine and VH and/or VL sequences are amplified from the patient and the antibody repertoire analysed as described above.

Qualitative and quantitative assessment of the vaccination process is possible:

- 1) The VH/VL repertoire of a group of patients who have been administered an experimental *C. difficile* vaccine is assessed. A precise molecular dissection of the resulting antibody response to the vaccine can be performed to determine (a) clonal expansion of protective antibodies, (b) protective antibody production in different populations (for example differing ethnic groups with different genetic backgrounds, as well as e.g. age and sex groups), and (c) the long term effect of the vaccine i.e. the antibody response over the long term – long term antibody memory in the immune system as well as any autoimmune defects;
- 2) Where vaccination results in an increased frequency of a given antibody, the antibody sequence can easily be cloned and expressed and used in animal models of infection. If the antibody is protective, it may be useful in itself as an immunotherapy, but this also demonstrates that the vaccine is likely to be protective without subjecting the human subject to experimental infection.

- 16 -

Also provided to the present invention is a method of manufacture of a medicament for the treatment of a *C. difficile* infection which produces at least one antigen, comprising the steps of:

- (i) performing a method according to the present invention to identify a set of candidate sequences for antibodies specific against the at least one antigen produced by *C. difficile*; and
- (ii) synthesising at least one antibody comprising a said candidate sequence specific against the at least one antigen produced by *C. difficile*.

Medicaments and methods of treatment according to the present invention will be readily apparent to one skilled in the art. Medicaments may be prepared using pharmaceutically acceptable carriers, diluents or excipients (Remington's: The Science and Practice of Pharmacy (1995) Mack Publishing Company, Easton, PA, USA). The medicaments and methods of treatment may be effected using a pharmaceutically effective amount of the antibody/antigen-binding fragment. Appropriate dosages will be readily apparent to one skilled in the art and may be readily determined, for example by means of dose-response experiments

Also provided according to the present invention is a method of treatment of an infection of a patient by *C. difficile* which produces at least one antigen, comprising the steps of:

- (i) performing a method according to the present invention to identify a set of candidate sequences for antibodies specific against the at least one antigen produced by *C. difficile*;
- (ii) synthesising at least one antibody comprising a said candidate sequence specific against the at least one antigen produced by *C. difficile*; and

- 17 -

- (iii) administering a therapeutically effective quantity of said at least one synthesised antibody to said patient.

Also provided according to the present invention is a method of producing a database which identifies candidate sequences for antibodies specific against at least one antigen produced by *C. difficile*, comprising the steps of:

- (i) performing a method according to the present invention to identify a set of candidate sequences for antibodies specific against the at least one antigen produced by *C. difficile*; and
- (ii) storing the data produced by said method in said database.

Also provided is a method of generating a report which identifies candidate sequences for antibodies specific against at least one antigen produced by *C. difficile*, comprising the steps of:

- (i) performing a method according to the present invention to identify a set of candidate sequences for antibodies specific against the at least one antigen produced by *C. difficile*; and
- (ii) producing a report comprising the data produced by said method of step (i).

Also provided according to the present invention is a method for determining the efficacy of a vaccine, comprising the steps of:

- (i) with B cells isolated from at least one patient who has been administered said vaccine, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells; and
- (ii) correlating said sequenced at least the CDR3 region of the VH and/or VL coding regions of said B cells to identify a set of candidate sequences for at least the CDR3 region of antibodies specific against

- 18 -

said vaccine, each of said set of CDR3 candidate sequences or a sequence having at least 80% homology therewith occurring in total at a frequency of at least 1 percent in the set of sequences determined at step (i).

5

In certain embodiments of the invention, step (i) above may comprise the steps of:

- (i)(a) administering said vaccine to at least one patient;
- (i)(b) isolating B cells from said at least one patient; and
- (i)(c) sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells.

10

Correlation step (ii) may comprise correlating said sequenced at least a CDR3 region of the VH and/or VL coding regions of said B cells with sequenced at least a CDR3 region of the VH and/or VL coding regions of B cells isolated from at least one patient who has been infected with *C. difficile* against which vaccination with said vaccine is intended to stimulate a protective immune response.

15

As described above, additional information may be used in the correlation of step (ii), including:

20

- (i) the time since administration of the vaccine to the patient;
- (ii) patient details - none or more of: sex, race, and age; and
- (iii) the range of complementarity determining regions (CDR) ie the variable regions of the antibody that undergo direct antigen binding/contact.

25

The antibody sequences determined from patients who have been administered the vaccine can be compared with antibody sequences isolated from patients who have been infected with *C. difficile* against which the vaccine is intended to stimulate a protective immune

- 19 -

response in patients. Thus it is possible to determine whether the vaccine results in the generation of antibodies which are also produced in response to infection by *C. difficile* itself. In particular, the comparison can be made using antibody sequences determined from patients who have recovered from infection by *C. difficile*.

5

In particular the method may determine the efficacy of the vaccine in stimulating a protective immune response against *C. difficile* against which vaccination with the vaccine is intended to stimulate a protective immune response.

10 Also provided is a method of producing a database which identifies the efficacy of a vaccine, comprising the steps of:

- (i) performing a method according to the present invention to determine the efficacy of said vaccine; and
- (ii) storing the data produced by said method in said database.

15

Also provided according to the present invention is a method of generating a report which identifies the efficacy of a vaccine, comprising the steps of:

- (i) performing a method according to the present invention to determine the efficacy of said vaccine; and
- 20 (ii) producing a report comprising the data produced by said method.

Also provided according to the present invention is a diagnostic test method for identifying a *Clostridium difficile* infection in a patient, comprising the steps of:

- (i) with B cells isolated from said patient, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells;
- 25 (ii) comparing said sequenced at least said CDR3 region of the VH and/or VL coding regions of said B cells with a set of sequences for at least the CDR3 region of antibodies specific against *Clostridium difficile*,

- 20 -

and determining whether each of said set of CDR3 sequences or a sequence having at least 80% homology therewith occurs in total at a frequency of at least 1 percent in the set of sequences determined at step (i); and

- 5 (iii) correlating the results of comparison step (ii) to determine the presence or absence of a *Clostridium difficile* infection in said patient.

Also provided according to the present invention is a diagnostic test method for determining the susceptibility of a patient to *Clostridium difficile* infection, comprising the steps of:

- 10 (i) with B cells isolated from said patient, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells;
- (ii) comparing said sequenced at least said CDR3 region of the VH and/or VL coding regions of said B cells with a set of sequences for at least the CDR3 region of antibodies specific against *Clostridium difficile*, and determining whether each of said set of CDR3 sequences or a sequence having at least 80% homology therewith occurs in total at a frequency of at least 1 percent in the set of sequences determined at step (i); and
- 15 (iii) correlating the results of comparison step (ii) to determine the susceptibility of said patient to *Clostridium difficile* infection.
- 20

The diagnostic test method could be used to sequence the CDR3 sequences of patients in for example a hospital ward, to determine which patients were susceptible to infection by the bacterium. The presence of *C. difficile* specific antibodies in a sample (as determined by sequencing of CDR3 regions) would likely indicate that such patients were capable of mounting a protective immune response toward the bacterium and would not require antibiotic treatment. In contrast, those patients identified who had not produced *C. difficile*

25

- 21 -

specific antibodies could be identified as being immunocompromised and therefore candidates for antibiotic treatment during the course of a *C. difficile* infection.

Also provided according to the present invention is a diagnostic test kit for performing a
5 diagnostic test method according to the present invention. The kit may contain reagents (e.g. PCR primers) which would enable a person skilled in the art to identify CDR3 sequences from a patient.

As regards antibodies usable in the present invention, particularly synthetic antibodies (the
10 term "antibody" is also considered to incorporate antigen binding fragments of antibodies unless the context otherwise requires), the antibody may be a whole antibody or an antigen binding fragment thereof and may in general belong to any immunoglobulin class. Thus, for example, it may be an IgM, IgG or an IgA antibody. The antibody or fragment may be of animal, for example, mammalian origin and may be for example of murine, rat, sheep
15 or human origin. It may be a natural antibody or a fragment thereof, or, if desired, a recombinant antibody fragment, i.e. an antibody or antibody fragment which has been produced using recombinant DNA techniques.

Particular recombinant antibodies or antibody fragments include, (1) those having an
20 antigen binding site at least part of which is derived from a different antibody, for example those in which the hypervariable or complementarity determining regions of one antibody have been grafted into the variable framework regions of a second, different antibody (as described in, for example, EP 239400); (2) recombinant antibodies or fragments wherein non-Fv sequences have been substituted by non-Fv sequences from other, different
25 antibodies (as described in, for example, EP 171496, EP 173494 and EP 194276); or (3) recombinant antibodies or fragments possessing substantially the structure of a natural immunoglobulin but wherein the hinge region has a different number of cysteine residues from that found in the natural immunoglobulin but wherein one or more cysteine residues

- 22 -

in a surface pocket of the recombinant antibody or fragment is in the place of another amino acid residue present in the natural immunoglobulin (as described in, for example, WO 89/01782 and WO 89/01974).

- 5 Teachings of texts such as Harlow, E. and Lane, D. ("Using Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, New York, 1998) further details antibodies, antibody fragments, their preparation and use.

10 The antibody or antibody fragment may be of polyclonal or monoclonal origin. It may be specific for at least one epitope.

Antigen binding antibody fragments include, for example, fragments derived by proteolytic cleavage of a whole antibody, such as F(ab')₂, Fab' or Fab fragments, or fragments obtained by recombinant DNA techniques, for example Fv fragments (as described, for
15 example, in WO 89/02465).

Where it is desired to produce recombinant antibodies according to the invention these may be produced using, for example, the methods described in EP 171469, EP 173494, EP 194276 and EP 239400. Antibody fragments may be produced using conventional
20 techniques, for example, by enzymatic digestion with pepsin or papain.

Antibodies according to the invention may be labelled with a detectable label or may be conjugated with an effector molecule, for example a drug eg. an antibacterial agent or a toxin or an enzyme, using conventional procedures and the invention extends to such
25 labelled antibodies or antibody conjugates.

The contents of each of the references discussed herein, including the references cited therein, are herein incorporated by reference in their entirety.

Where "PMID:" reference numbers are given for publications, these are the PubMed identification numbers allocated to them by the US National Library of Medicine, from which full bibliographic information and abstract for each publication is available at www.ncbi.nlm.nih.gov.

5

The invention will be further apparent from the following description, with reference to the several figures of the accompanying drawings, which show, by way of example only, forms of identifying candidate sequences for antibodies specific against at least one antigen and of determining the efficacy of a vaccine.

10

Of the Figures:

15

Figure 1 shows the general principles for the isolation and DNA sequencing of VH and/or VL antibody gene fragments from B cells. Reference numeral 10 indicates primary PCR, reference numeral 20 the cloning into a DNA sequencing vector using the T-tailing principle - random orientation, and reference number 30 the determining of the nucleotide sequence of forward and reverse strands using M13 forward and reverse primers; and

20

25

Figure 2 shows a schematic depiction of resynthesised recombinant antibody gene cassette. VH and VL regions are linked with a glycine serine-rich linker. Each variable domain contains three Complementarity Determining Regions (CDRs) which participate in antigen binding.

- 24 -

EXPERIMENTS

The experiments below detail methods for identifying antibody VH and/or VL CDR sequences that confer immunity in patients with *C. difficile* infection. These sequences can then be used to produce synthetic antibodies and these synthetic antibodies are suitable for administration to patients for therapy. Also detailed are the VH and VL CDR3 sequences identified by the methods.

Also detailed are methods of determining the efficacy of a vaccine in generating a desired immune response in a patient.

Unless stated otherwise, all procedures were performed using standard protocols and following manufacturer's instructions where applicable. Standard protocols for various techniques including PCR, molecular cloning, manipulation and sequencing, the manufacture of antibodies, epitope mapping and mimotope design, cell culturing and phage display, are described in texts such as McPherson, M.J. *et al.* (1991, PCR: A practical approach, Oxford University Press, Oxford), Sambrook, J. *et al.* (1989, Molecular cloning: a laboratory manual, Cold Spring Harbour Laboratory, New York), Huynh and Davies (1985, "DNA Cloning Vol I - A Practical Approach", IRL Press, Oxford, Ed. D.M. Glover), Sanger, F. *et al.* (1977, PNAS USA 74(12): 5463-5467), Harlow, E. and Lane, D. ("Using Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, New York, 1998), Jung, G. and Beck-Sickinger, A.G. (1992, Angew. Chem. Int. Ed. Eng., 31: 367-486), Harris, M.A. and Rae, I.F. ("General Techniques of Cell Culture", 1997, Cambridge University Press, ISBN 0521 573645), "Phage Display of Peptides and Proteins: A Laboratory Manual" (Eds. Kay, B.K., Winter, J., and McCafferty, J., Academic Press Inc., 1996, ISBN 0-12-402380-0).

- 25 -

Reagents and equipment useful in, amongst others, the methods detailed herein are available from the likes of Amersham (www.amersham.co.uk), Boehringer Mannheim (www.boehringer-ingenelheim.com), Clontech (www.clontech.com), Genosys (www.genosys.com), Millipore (www.millipore.com), Novagen (www.novagen.com),
5 Perkin Elmer (www.perkinelmer.com), Pharmacia (www.pharmacia.com), Promega (www.promega.com), Qiagen (www.qiagen.com), Sigma (www.sigma-aldrich.com) and Stratagene (www.stratagene.com).

The term "antibody" in its various grammatical forms is used herein to refer to
10 immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antibody combining site or paratope. Such molecules are also referred to as "antigen binding fragments" of immunoglobulin molecules.

15 Illustrative antibody molecules are intact immunoglobulin molecules, substantially intact immunoglobulin molecules and those portions of an immunoglobulin molecule that contain the paratope, including those portions known in the art as Fab, Fab', F(ab')₂, scFv and F(v).

The term "antibody combining site" refers to that structural portion of an antibody
20 molecule comprised of a heavy and light chain variable and hypervariable regions that specifically binds (immunoreacts with) antigen.

The identification and sequencing of B cells producing antibodies and the analysis of the resulting data comprises the following basic steps:

- 25
- (1) Isolation of VH and/or VL coding regions from circulatory B cells of human patients.
 - (2) Determining the Nucleotide sequence of VH and/or VL repertoires.
 - (3) Determination of VH and/or VL primary amino acid sequences.

- 26 -

- (4) Extraction of CDR regions *in silico* - incorporation into the database
- (5) Detection of dominant CDR & framework regions in VH and/or VL repertoire.
- (6) Construction & Production of therapeutic recombinant antibodies from dominant antibody sequences.

1. Isolation of VH and/or VL coding regions from circulatory B cells of human patients

For this method, human patients with *C. difficile* infection were selected as donors of immunised B cells. The criteria for selection were:

(a) The patient must exhibit a pronounced antibody response to *C. difficile*, detectable for example by Western blotting. Samples of antibodies were collected from a patient blood sample. Samples were diluted and tested for immunoreactivity by Western blotting using antigen(s) derived from *C. difficile*. In such an assay, patients exhibiting strong antibody responses showed pronounced recognition/antibody binding of *C. difficile* antigens as detected using anti-human polyclonal detection antibodies.

(b) The patient has survived the course of *C. difficile* infection - this improves the chance of generating B cell responses for antibodies capable of neutralizing the bacterium.

Peripheral B-cell lymphocytes (PBLs) were collected from infected patient blood samples. For this heparinised blood was diluted in PBS (20 ml total) and overlaid onto a 15 ml cushion of Ficol Hypaque (Pharmacia; unless stated otherwise, all chemicals and culture media were purchased from Sigma, UK) in a 30 ml centrifuge tube. The PBLs were then collected by centrifugation (400 x g, 5 minutes) and washed in PBS and harvested by centrifugation again. RNA was prepared from PBLs using the QuickPrep mRNA purification kit (Pharmacia) exactly according to the manufacturers instructions. The

- 27 -

isolated mRNA was used to prepare cDNA via a reverse transcriptase reaction (Promega cDNA synthesis kit). For this 2 µg of mRNA was re-suspended in 16 µL nuclease-free water and heated to 65 °C for 3 minutes (to denature secondary structure) and then immediately chilled on ice for 1 minute. The mRNA was then added to the following cocktail: 8 µL 25 mM MgCl₂, 4 µL dNTP mix (10 mM with respect to each ribonucleotide triphosphate), 1 µL RNAsin 40 u µL⁻¹ stock solution, 1.2 µL AMV reverse transcriptase (25 u µL⁻¹ stock solution), 6 µL cDNA 10 pmol µL⁻¹ primer (see Figure 1 - cDNA synthesis). The mixture was incubated at 42 °C for 1 hour and then incubated at 100 °C for 3 minutes to stop the reaction.

DNA coding for antigen binding regions was then amplified by PCR using the cDNA prepared from patients PBLs. For this, the cDNA was used in the following PCR reaction to produce either heavy chain or light chain-derived antibody variable region DNA (see Figure 1): 2.5 µL cDNA, 33 µL water, 4 µL dNTP mix (25 mM with respect to each deoxynucleotide triphosphate), 5 µL Taq reaction buffer (Perkin Elmer), 2.5 µL of an equimolar primer mix (final concentration of 20 pmol with respect to each primer) and 0.5 µL Taq DNA polymerase (1 u µL⁻¹, Perkin Elmer Corp). The forward and reverse primers used in these reactions are depicted in Figure 1, and their respective nucleotide sequences are listed in Table 1. PCR reaction conditions were 94 °C for 1 minute, 57 °C for 1 minute and 72 °C for 2 minutes for a total of 30 cycles, with an extended denaturation (94 °C for 5 minutes) prior to cycle 1 and an additional extension step after the end of cycle 30 (72 °C for 10 minutes). PCR was performed using a Perkin Elmer 9700 GeneAmp PCR machine. 5 µL of the PCR reaction was run on a 1% agarose gel to check the amplification of the expected 393 base pair (bp) product. The remaining product was run on a 0.8% low melting point (LMP) agarose gel and the 393 bp band was excised using a clean scalpel blade. DNA was extracted from the agarose gel slice using a GeneClean II Kit (Anachem, Luton, UK).

- 28 -

The PCR resulted in two fragment types –derived from VH and VL regions respectively, depending on the PCR primer sets used (for details of primers sets for antibody genes and a PCR schematic see Table 1 and Figure 1, respectively).

Table 1: Primers used in generating and sequencing VH and VL gene fragments. All sequences are given in the 5' to 3' direction. Use of primers is described above and in Figure 1.

Table 1

<u>Primers used in cDNA synthesis reactions</u>									SEQ ID NO:
HuIgG1 - 4FOR	GTC	CAC	CTT	GGT	GTT	GCT	GGG	CTT	01
HuCLFOR	AGA	CTC	TCC	CCT	GTT	GAA	GCT	CTT	02
<u>Primers for primary PCR</u>									SEQ ID NO:
HuJH1 - 2FOR	TGA	GGA	GAC	GGT	GAC	CAG	GGT	GCC	03
HuJH3FOR	TGA	AGA	GAC	GGT	GAC	CAT	TGT	CCC	04
HuJH4 - 5FOR	TGA	GGA	GAC	GGT	GAC	CAG	GGT	TCC	05
HuJH6FOR	TGA	GGA	GAC	GGT	GAC	CGT	GGT	CCC	06
HuVH1aBACK	CAG	GTG	CAG	CTG	GTG	CAG	TCT	GG	07
HuVH2bBACK	CAG	GTG	AAC	TTA	AGG	GAG	TCT	GG	08
HuVH3aBACK	CAG	GTG	CAG	CTG	GTG	GAG	TCT	GG	09
HuVH4aBACK	CAG	GTG	CAG	CTG	CAG	GAG	TCG	GG	10
HuVH4bBACK	CAG	GTG	CAG	CTA	CAG	CAG	TGG	GG	11
HuVH5aBACK	GAG	GTG	CAG	CTG	TTG	CAG	TCT	GC	12
HuVH6aBACK	CAG	GTA	CAG	CTG	CAG	CAG	TCA	GG	13
HuJK1FOR	ACG	TTT	GAT	TTC	CAC	CTT	GGT	CCC	14
HuJK2FOR	ACG	TTT	GAT	CTC	CAG	CTT	GGT	CCC	15
HuJK3FOR	ACG	TTT	GAT	ATC	CAC	TTT	GGT	CCC	16
HuJK4FOR	ACG	TTT	GAT	CTC	CAC	CTT	GGT	CCC	17
HuJK5FOR	ACG	TTT	ATT	CTC	CAG	TCG	TGT	CCC	18
HuVK1BACK	GAC	ATC	CAG	ATG	ACC	CAG	TCT	CC	19
HuVK2BACK	GAT	GTT	GTG	ATG	ACT	CAG	TCT	CC	20
HuVK3BACK	GAA	ATT	GTG	TTG	ACG	CAG	TCT	CC	21
HuVK4BACK	GAC	ATC	GTG	ATG	ACC	CAG	TCT	CC	22
HuVK5BACK	GAA	ACT	ACA	CTC	ACG	CAG	TCT	CC	23
HuVK6BACK	GAA	ATT	GTG	CTG	ACT	CAG	TCT	CC	24
<u>Primers for DNA Sequencing</u>									
M13forward	GTA	AAA	CGA	CGG	CCA	GT			25
M13reverse	AAC	AGC	TAT	GAC	CAT	G			26

- 29 -

VH and/or VL gene fragments were cloned into cloning vector pGEM-T easy (Promega Corporation) to facilitate DNA sequencing. For this 3 µg PCR product was prepared for restriction using QIAquick PCR purification spin columns (Qiagen, UK) according to the manufacturers instructions. DNA was eluted from the spin column in 40 µL buffer EB.

- 5 Purified PCR product (2 µL) was mixed with 1 µL pGEM-T easy vector, 6 µL water and 1 µL DNA ligase and the mixture ligated for 1 h at room temperature. Ligations were then transformed into electrocompetent *E. coli* TG1 cells (Stratagene) by electroporation, and plated out onto agar plates containing Ampicillin 100 µg ml⁻¹ IPTG (100 µM) and X-gal (0.006 % w/v). Colonies were allowed to grow overnight at 37 °C and then stored at 4 °C.
- 10 Recombinant colonies are identified as white colonies on this media.

2. Determining the nucleotide sequence of VH and/or VL repertoires

- DNA was first prepared from VH and/or VL recombinant *E. coli* clones. For this, individual colonies were each transferred to 1.2 ml LB broth supplemented with
- 15 Ampicillin 100 µg ml⁻¹ using a 96 well plate format. Cultures were the grown at 37 °C for 24 hours. Bacterial cells were harvested by centrifugation at 4,000 x g, 30 minutes and the supernatants discarded. Plasmid DNA was prepared using Wizard SV 96 plasmid purification kits (Promega Corporation) essentially following the manufacturer's instructions. Yields of plasmid DNA were typically in the order of 5 µg per 1.2 ml starter
- 20 culture.

- DNA sequencing reactions were performed using the DYEnamic ET dye terminator cycle sequencing kit (Amersham Pharmacia Biotech). Purified plasmid DNA (0.5 µg) was mixed with 8 µL DYEnamic ET terminator reagent premix and 1 µL M13 forward or reverse
- 25 primer (5 µM) in a total reaction volume of 20 µL. Thermal cycling was then performed using a GeneAmp PCR system 9700 (Perkin Elmer) with the following parameters: 95 °C, 20 seconds; 50 °C, 15 seconds; 60 °C, 1 minute; 30 cycles). Reactions were performed using 96 well format non-skirted ELISA plates (AB Gene). Unincorporated dye terminator

- 30 -

were removed using precipitation. For this, ethanol samples were mixed with 2 μ L 7.5 M ammonium acetate and 55 μ L of 100 % ethanol and centrifuged at 3000 g for 30 minutes. DNA pellets were washed with 70 % ethanol and re-suspended in 20 μ L loading solution. Reactions were sequenced using a MegaBACE 1000 DNA sequencer (Amersham Pharmacia Biotech) following the manufacturers instructions (2 kV injection voltage for 30 s with electrophoresis at 6 kV for 200 minutes). Chromatograms were exported using the .scf file format for finishing and archiving.

3. Determination of VH and/or VL primary amino acid sequences

DNA sequences of VH and/or VL were determined in both forward and reverse strands (using M13 forward and reverse primers respectively) and compared in order to highlight discrepancies and maintain a high degree of accuracy. This was done using the Staden suite of programs (Staden, R. (1996) *The Staden Sequence Analysis Package*. Molecular Biotechnology 5, 233-241). First, sequence .scf files were entered into the PREGAP program where vector sequence was stripped off and the quality of the sequence was assessed. Poor quality sequences where the DNA sequence was ambiguous were rejected and re-sequenced. Forward and reverse strands were matched using the GAP program to highlight and resolve areas containing sequencing artefacts. The orientation of the VH and/or VL sequence was noted and reverse complemented where necessary to produce only forward reading frame orientations for translation. VH and VL gene sequences were then translated into amino acid sequences. Each sequence was given a unique identifier (name).

4. Extraction of CDR regions *in silico* - incorporation into the database

A general teaching of identifying CDR regions is at www.bioinf.org.uk/abs/

The following set of rules will allow the definition of the CDRs in an antibody sequence. There are rare examples where these virtually constant features do not occur (for example

- 31 -

the human heavy chain sequence EU does not have Trp-Gly after CDR-H3). The Cys residues are the best-conserved feature.

CDR-L1

5	Start	Approx. residue 24
	Residue before	always a Cys
	Residue after	always a Trp. Typically Trp-Tyr-Gln, but also, Trp-Leu-Gln, Trp-Phe-Gln, Trp-Tyr-Leu
	Length	10 to 17 residues

10

CDR-L2

	Start	always 16 residues after the end of CDR-L1
	Residues before	generally Ile-Tyr, but also, Val-Tyr, Ile-Lys, Ile-Phe
	Length	always 7 residues

15

CDR-L3

	Start	always 33 residues after end of CDR-L2
	Residue before	always Cys
	Residues after	always Phe-Gly-Xaa-Gly (SEQ ID NO: 92)
20	Length	7 to 11 residues

CDR-H1

	Start	Approx. residue 26 (always 4 after a Cys)
	Residues before	always Cys-Xaa-Xaa-Xaa (SEQ ID NO: 93)
25	Residues after	always a Trp. Typically Trp-Val, but also, Trp-Ile, Trp-Ala
	Length	10 to 12 residues

CDR-H2

- 32 -

	Start	always 15 residues after the end of CDR-H1
	Residues before	typically Leu-Glu-Trp-Ile-Gly (SEQ ID NO: 94), but a number of variations
	Residues after	Lys/Arg-Leu/Ile/Val/Phe/Thr/Ala-Thr/Ser/Ile/Ala
5	Length	16 to 19 residues;

CDR-H3

	Start	always 33 residues after end of CDR-H2 (always 2 after a Cys)
	Residues before	always Cys-Xaa-Xaa (typically Cys-Ala-Arg)
10	Residues after	always Trp-Gly-Xaa-Gly (SEQ ID NO: 95)
	Length	3 to 25 residues

5. Detection of dominant CDR & framework regions in VH and/or VL repertoire

The database was constructed using SQL Server Database software (Microsoft). This
15 allowed the database to be queried using SQL and allowed CDR1, CDR2 and CDR3 sequences to be extracted from any range of database VH and/or VL sequences in FASTA format. Extracted CDRs were then subject to multiple alignment using CLUSTAL X in such a way so that identical or very similar CDRs are grouped together in blocks (Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) CLUSTAL W: improving the
20 sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. Nucleic Acids Research, 22:4673-4680). From this the frequency of recurring CDRs was determined and CDR-dominance in the immunoglobulin repertoire was established. Alternatively, a graphical interface was employed using Probiosys (proBionic EMEA, Amsterdam). ProBiosys
25 prepares dendrograms from CLUSTAL .phy files and allows a rapid visual interpretation of the relationships among large numbers of aligned sequences. The whole process was performed for CDRs from the same patient sampled at different points during an illness, for CDRs from patients with different infections, or from sex-matched, age-matched or

race-matched patient groups, depending on the search and the additional information added to the database entries.

5

6. Construction & production of therapeutic recombinant antibodies from dominant antibody sequences

Once dominant antibody sequences have been recognised in a given repertoire, the information can be used to infer the presence of CDRs and frameworks that confer
10 immunity. Selected VH and/or VL sequences can be identified and their gene sequences resynthesised using synthetic oligonucleotides. This is important as it also allows the human-derived gene sequence to be codon-optimised for *E. coli* in order to improve protein expression. Dominant VH and/or VL sequences can be spliced together using a spacer (linker) sequence (Figure 2). This gene cassette, termed a scFv, can be resynthesised
15 to include terminal *NdeI* and *NotI* restriction sites for cloning into expression vector pET29b (Novagen). ScFv DNA can be cut with *NdeI* (cuts in VH) or *NotI* (cuts in VL) using the following reaction: 40 µl DNA (4 µg), 10 µl of Restriction enzyme buffer D (Promega), 47 µl water and 2 µl *NdeI* and *NotI* enzyme (10 u µl⁻¹; Promega) with digestion for 4 h at 37 °C. DNA can be then fractionated on 0.7% agarose TAE gels and
20 the digested DNA excised from the gel and purified from the agarose slice using the GeneClean II kit (Bio 101) exactly according to the manufacturers instruction. pET29b vector DNA can be cut with *NdeI* and *NotI* as described above. 1 µg vector can be mixed with 1 µg restricted VL DNA resuspended in 8.5 µl water and ligated by addition of 1 µl 10 × ligation buffer (Boehringer Mannheim) and 0.5 µl DNA ligase (3 u µl⁻¹; Boehringer
25 Mannheim), followed by ligation overnight at 14 °C.

The entire ligation can be transformed into *E. coli* TG2 cells (Stratagene, UK) by electroporation. Transformants can be plated out onto agar plates containing Tetracycline

- 34 -

(25 $\mu\text{g ml}^{-1}$ - selective for *recA* ::Tn10 on the chromosome of TG2) and Kanamycin (50 $\mu\text{g ml}^{-1}$ - selective for pET29). Recombinant plasmid DNA prepared from individual clones can be checked for scFv sequence by digestion with *NdeI* and *NotI* as described above.

5

Recombinant scFvs can be over-expressed in *E. coli* strain JM109(DE3) and purified thus enabling biochemical and biological characterisation. Recombinants can be spread on LB plates supplemented with 50 $\mu\text{g ml}^{-1}$ kanamycin and a single colony can be used to inoculate a 10 microlitre starter culture of LB broth with 50 $\mu\text{g ml}^{-1}$ kanamycin. For this, a 1 L expression culture can be prepared in the presence of kanamycin for 4-5 h at 37 °C with shaking at 300 rpm. At $\text{OD}_{600}=1$, induction can be performed using IPTG and the cells can be grown with vigorous aeration for a further 3-5 h. Cells can then be harvested by centrifugation (4000 x g, 15 minutes, 4 °C). The cell pellet can be resuspended in 10 ml fresh (4 °C) lysis buffer (below) and stored at -20 °C overnight prior to cell breakage using the 25 ml X-Press system (AB Biox)

10

15

LYSIS BUFFER

50 mM Tris HCl pH 8.0

1 mM EDTA

20 100 mM KCl

0.1 mM AEBSF (amino ethyl benzene sulphonic acid)

The lysate can be centrifuged in an Oakridge tube (24 300 x g, 4 °C, 30 minutes). The pellet can be resuspended in 20 ml of ice cold lysis buffer on ice using a Silverson lab blender. The lysate can be split into 8 x Oakridge tubes and each aliquot can be diluted to 30 ml (total 120 ml) in ice-cold Lysis Buffer. Inclusion bodies can be pelleted (24 300 x g, 4 °C, 30 minutes) and the supernatant discarded. Each of the 8 pellets can be resuspended in 30 ml of lysis buffer, and the inclusion bodies can be harvested by

25

- 35 -

centrifugation again (24 300 x g, 4 °C, 30 minutes). This wash/centrifugation step can be performed five times in total to clean the inclusion body fraction. Each pellet can be resuspended in 15 ml of ice-cold water and the inclusion bodies can be stored at -20 °C.

- 5 For refolding, 200 ml solution of 2% (w/v) N-lauryl sarcosine (NLS) in 50 mM Tris HCl pH 9.0 can be prepared as follows. 4 g NLS can be solubilised in 100 ml water with stirring. 10 ml of 1M Tris pH 9.0 stock solution can be added and made up to 195 ml with water. 5 ml of inclusion body slurry can be added and stirred vigorously for 30 minutes at room temperature.

10

CuCl₂ can be added to a concentration of 100 µM. This serves as a catalyst for oxidation. The refolding reaction can be transferred to 4 °C and stirred vigorously for 2 days to promote aeration. The refolding reaction can be vacuum filtered through a 0.44 µM vacucup bottle top filter unit (90 mm diameter, Gellman Sciences) and the filtrate
15 transferred to a Pellicon Labscale TFF system fitted with PLGC10 membrane unit (Millipore). The reaction can be concentrated to 25 ml using tangential flow, discarding the permeate (the scFv is localised to the retentate). The solution can be diafiltered against 40 x turn-over volumes (1 L) of 10 mM ammonium acetate (AAT) pH 9.0. Finally, the volume of the antibody can be diluted to 50 ml using 10 mM AAT pH 9.0. The buffer
20 exchanged antibody can be stored for 2 hours at 4 °C. The typical protein content was 1 - 2 mg ml⁻¹ with a yield of up to 50 mg per litre.

25

For scFv purification a 10 ml bed volume of Ni NTA superflow agarose Qiagen can be prepared according to the manufacturers instructions using a 10 ml glass column (Sigma) without flow adaptors. The column can be equilibrated with 50 ml Buffer B (6M urea, 0.1 M NaH₂PO₄, 10 mM Tris HCl pH 8.0). Refolded scFv (as described above) can be diluted 1/5 in Buffer B and applied directly to the column at ml min⁻¹. 50 ml buffer B can be applied to the column (flow rate 5 ml min⁻¹) followed by 50 ml Buffer C (same

- 36 -

composition as buffer B but pH 6.3). The purified scFv can be eluted from the column by slowly applying Buffer B supplemented with 250 mM high grade imidazole until the $A_{280} < 0.05$.

5 Results

The variable heavy chain (VH) and variable light chains of the antibody genes from peripheral blood lymphocytes (PBLs) isolated from *C. difficile* infected patients have been TA cloned and sequenced. In each case the third complementarity determining region (i.e. CDR-H3 or CDR-L3) was identified. This region is both the most variable region of the
10 VH chain and the area identified as being most important in determining antigen binding, for this reason this area is used as the signature for each VH chain.

Clinical Information

Patient D01 - Variable heavy chains (VH's) and variable light chains (VL's) were
15 sequenced from PBLs taken ten days after the onset of CDAD. Lab results confirmed *C. difficile* toxin was present.

Patient D02 - VH's and VL's were sequenced from PBLs taken ten days after the onset of CDAD. Lab results confirmed *C. difficile* toxin was present. In the seven subsequent
20 weeks after sample was taken, the patient did not develop any further *C. difficile* infections, suggesting that the antibodies produced in response to the infection were protective.

Patient D03 - VH's and VL's were sequenced from PBLs taken ten days after the onset of
25 CDAD. Lab results confirmed *C. difficile* toxin was present. In the three subsequent weeks after sample was taken, the patient did not develop any further *C. difficile* infections, suggesting that the antibodies produced in response to the infection were protective.

- 37 -

Sequencing results

Patient D01 showed a high degree of focusing with both VH's and VL's. Three different CDR3 sequences make-up over 80 % of the sequenced VH's. VH's with the CDR3 sequence of SEQ ID NO: 27 account for 57.8 % of sequenced VH's from this patient. 5 VH's with the CDR3 sequence of SEQ ID NO: 28 account for 13.5 % of all VH's, and VH's the CDR3 sequence of SEQ ID NO: 29 account for 10 % of all sequenced VH's in this patient. Three different CDR3 sequences also make up the majority of sequenced VL's, these are the CDR3 sequences of SEQ ID NOs: 32, 34, and 33 (occurring at frequencies of 32.7 %, 15.3%, and 6.6%, respectively).

10

Similarity searches were performed using these common VH and/or VL sequences against other VH and/or VL sequences contained within a database (FABTEC database). which contains VH and/or VL regions of PBLs isolated from patients with methicillin-resistant *Staphylococcus aureus* (MRSA), *Candida albicans*, *Pseudomonas aeruginosa* (the 15 causative agent of cystic fibrosis), *Streptococcus oralis* and vancomycin-resistant Enterococci (VRE) infections and to identify antibody sequences which are associated with resistance to the infections and which can therefore be used to effect therapy against those infections. These sequences have formed the basis of WO 01/76627. For all of these, the isolated VH and/or VL sequences demonstrate skewing of the antibody repertoire over the 20 course of infection, thus revealing the identity of matured, immunity-conferring VH and/or VL sequences. This has typically been done using 5,000 to 15,000 sequences for each VH and/or VL. VH and/or VL repertoires from healthy individuals can also be used for comparison purposes.

20

25 Using the BLOSUM50 matrix it was possible to pick out groups of antibodies with CDR3's that had greater than 70% similarity. The CDR3 sequences of SEQ ID NOs: 27 and 28 (Table 2 and 3, respectively), showed sequence similarity of greater than 70% only

- 38 -

with other VH's from *C. difficile* infected patients VH's sequenced from individuals with other infections (*C. albicans*, MRSA and VRE) showed no similar CDR3 sequences.

In the following Tables, libraries starting D are derived from *C. difficile* infected patients, libraries starting M are derived from MRSA infected patients, libraries starting V are derived from VRE infected patients; and libraries starting C are derived from *C. albicans* infected patients.

Table 2

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
27	100.0	D01	184
27	100.0	D03	5
35	100.0	D01	1
36	100.0	D01	3
37	100.0	D01	1
37	100.0	D03	1
38	93.75	D01	1

Table 3

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
28	100.0	D01	43
28	100.0	D03	1
39	100.0	D01	1
40	88.88	D01	1

VH's with the CDR3 sequence of SEQ ID NO: 29 showed similarity to VH's from other *C. difficile* infected patients and MRSA infected patients (Table 4).

Table 4

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
41	100.0	D01	1
29	100.0	D01	32
29	100.0	M01	1
29	100.0	M02	5
29	100.0	M03	2
29	100.0	M04	8
42	100.0	M03	1
43	100.0	D01	1
44	93.75	D01	1
45	93.75	D01	1
46	75.0	M04	1
47	70.58	M05	1

The CDR3 sequences from the VL's of patient D01 have also had a similarity search performed on them. VL's with the CDR3 sequences of SEQ ID NOs: 32 and 33 (Tables 5 and 6, respectively) show a high degree of homology with CDR3 sequences from *C. difficile* infected patients but no homology to CDR3 sequences derived from patients with different clinical infections. VL's with the CDR3 sequence of SEQ ID NO: 34 showed similarity to VL's isolated from *C. difficile*, MRSA and VRE infected patients (Table 7).

- 40 -

Table 5

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
32	100.0	D51	64
48	90.9	D52	2
49	90.9	D52	1
50	90.9	D52	1
51	90.0	D51	1
52	90.0	D52	1
53	81.8	D52	1
54	81.8	D52	1
55	81.8	D52	2
56	81.8	D52	1
57	80.0	D52	1
58	72.7	D52	1

Table 6

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
33	100.0	D51	13
59	90.9	D52	1

- 41 -

Table 6 continued

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
60	83.3	D52	1
61	81.8	D52	4
62	81.8	D52	1
63	81.8	D52	1
64	81.8	D52	1
65	81.8	D51	1
66	81.8	D52	1
67	81.8	D52	1
68	72.7	D52	1
69	72.7	D52	1

Table 7

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
34	100.0	D51	30
34	100.0	M56	2
34	100.0	V54	2
70	88.9	D51	1
71	88.9	D51	1
72	88.9	V54	2
73	77.9	M55	1
74	77.9	D51	1
75	77.9	M55	1
76	77.9	M55	1

- 42 -

Patient D02 also showed a high degree of focusing of the VH repertoire, although focusing is less obvious in the VL repertoire. Two types predominate in the VH library of this patient. VH's with the CDR3 sequence of SEQ ID NO: 30 accounted for 45.6% of sequenced clones and those with the CDR3 sequence of SEQ ID NO: 31 accounted for 8.8% of the library. In the VL library the three most frequently isolated clones with the CDR3 sequences of SEQ ID NOs: 61, 96, and 97 were found at 4%, 4% and 3%, respectively.

Similarity searches in the FABTEC database showed the two common VH CDR3 sequences to be specific to *C. difficile* patients (Tables 8 and 9). VH's with the CDR3 sequence of SEQ ID NO: 30 match with other antibodies in that library and antibodies from library D03 (Table 8). VH's with the CDR3 sequence of SEQ ID NO: 31 match with antibodies from library D03 (Table 9).

Table 8

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
77	100.0	D02	1
78	100.0	D02	1
30	100.0	D02	52
30	100.0	D03	1
79	100.0	D02	1
79	100.0	D03	1
80	94.4	D02	1
81	94.4	D02	1
82	94.4	D02	1
83	88.8	D02	1
84	77.7	D03	1

Table 9

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
31	100.0	D02	10
31	100.0	D03	4
85	90.0	D03	1
86	90.0	D03	1

Similarity searches using some of the VL's showed a clustering of sequences. 15% of VL's from patient D02 (library D52) showed high degree of similarity to the CDR3 sequence of SEQ ID NO: 48, VL's from patient D01 (library D51) also showed a high degree of homology to this sequence (Table 10). In a similar way a cluster around the sequence of SEQ ID NO: 61 accounts for 13% of patient D02 sequenced VL's, again clones similar to this one are also found in patient D01 (Table 11). Neither of these clones show any significant similarity to VL CDR3 sequences isolated from patients infected with other pathogens. It should also be note that using the BLOSUM50 matrix these two clones show greater than 60% similarity to each other.

Table 10

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
48	100.0	D52	2
49	100.0	D52	1
53	100.0	D52	1
32	90.9	D51	64
87	81.8	D52	2
51	81.8	D51	1
88	81.8	D52	1

- 44 -

Table 10 continued

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
57	72.7	D52	1
89	72.7	D52	1
50	72.7	D52	1
54	72.7	D52	1
55	72.7	D52	2

Table 11

CDR3 SEQ ID NO:	Similarity	Library	No. of clones
61	100.0	D52	4
65	100.0	D51	1
59	81.8	D52	1
33	81.8	D51	13
62	81.8	D52	1
63	81.8	D52	1
60	75.0	D52	1
64	72.7	D52	1
66	72.7	D52	1
68	72.7	D52	1
67	72.7	D52	1

Patient D03 has only preliminary results for VH's but shows many similar sequences to both D01 and D02 so are worth reporting. Only data from 49 VH sequences have been produced. 10% of the clones have the CDR3 sequence of SEQ ID NO: 27, this is the sequence of the most common VH in patient D01. 8% of VH's have the CDR3 sequence of SEQ ID NO: 31 - this is the second most common VH CDR3 sequence in patient D02.

- 45 -

Cross-Library comparison (Table 12) of the CDR3 sequences of VH's has shown 8 CDR3 sequences which appear in more than one of the *C. difficile* infected patients. The CDR3 sequences of SEQ ID NOs: 27, 28, and 37 appear in patients D01 and D03 and the sequences of SEQ ID NOs: 31, 90, 91, 79, and 30 all appear in patients D02 and D03.

5 So far patients D01 and D02 show no common CDR3 sequences. VL's have only been sequenced from patients D01 and D02, these patients share no identical CDR3 sequences.

Table 12

VH CDR3 SEQ ID NO:	Patient
27	D01 and D03
28	D01 and D03
37	D01 and D03
31	D02 and D03
90	D02 and D03
91	D02 and D03
79	D02 and D03
30	D02 and D03

Summary

The presence of a large number of a particular VH CDR3 sequence in a patient with a *C. difficile* infection indicates that VH as part of an scFv may be protective against the organism. CDR3 sequences that are shared by more than one patient indicate specific *C. difficile* sequences. It should also be noted that sequences with a high degree of homology often due to somatic mutation within the patient may also represent important sequences with similar but subtly different properties. Analysis of the CDR3 sequences of the VH from our individuals has lead to the identification of 17

- 46 -

potentially protective VH's. These VH CDR3 sequences correspond to SEQ ID NOs: 30, 79, 28, 91, 41, 39, 38, 27, 36, 36, 40, 37, 90, 29, 44, 45, and 43.

5 A similar analysis of the VL has identified 44 different VL CDR3 sequences. The larger number is due to the increased rate of somatic mutation in antibody light chains leading to VL's with slightly different sequences. These VL CDR3 sequences correspond to SEQ ID NOs: 59, 60, 61, 33, 62, 63, 64, 65, 66, 77, 84, 78, 30, 79, 80, 81, 82, 87, 57, 74, 85, 31, 86, 82, 51, 48, 88, 49, 53, 32, 52, 70, 71, 34, 89, 68, 67, 50, 54, 55, 56, 59, 58, 96, and 97.

- 47 -

CLAIMS

1. An antibody or an antigen binding fragment thereof having the CDR-H3 sequence selected from the group consisting of: SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, and SEQ ID NO: 31.

2. An antibody or an antigen binding fragment thereof having the CDR-L3 sequence selected from the group consisting of: SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO: 34.

3. An antibody or an antigen binding fragment thereof having a CDR-H3 sequence selected from the group consisting of: SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 29, SEQ ID NO: 30, and SEQ ID NO: 31, and a CDR-L3 sequence selected from the group consisting of: SEQ ID NO: 32, SEQ ID NO: 33, and SEQ ID NO: 34.

4. A method for identifying candidate sequences of at least the CDR3 region of antibodies specific against at least one antigen produced by *Clostridium difficile* during an infection or against a vaccine, comprising the steps of:

- (i) with B cells isolated from at least one patient who has been infected by *Clostridium difficile* or administered said vaccine, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells; and
- (ii) correlating said sequenced at least the CDR3 regions of the VH and/or VL coding regions of said B cells from said at least one patient to identify a set of candidate sequences for at least a CDR3 region of antibodies specific against said at least one antigen produced by *Clostridium difficile* or against said vaccine, each of said set of candidate CDR3 sequences or a sequence having at least 80% homology therewith occurring in total at a frequency of at least 1 percent in the set of sequences determined at step (i).

- 48 -

5. A method according to claim 4, said B cells being selected from the group consisting of peripheral B-cell lymphocytes and B cells from the spleen.

6. A method according to claim 5, said peripheral B-cell lymphocytes being isolated
5 from blood from said at least one patient.

7. A method according to any of claims 4-6, said at least one antigen being an immunogen.

10 8. A method according to any of claims 4-7, said at least one patient displaying a pronounced antibody response in response to infection by *Clostridium difficile*.

9. A method according to any of claims 4-8, said at least one patient having recovered from infection by *Clostridium difficile*.

15 10. A method according to any of claims 4-9, said correlation step (ii) comprising determining putative amino acid sequences from said sequenced at least the VH and/or VL CDR3 coding regions, and correlating said putative amino acid sequences.

20 11. A method according to claim 9, said correlation step (ii) comprising identifying the Complementarity Determining Regions comprised in said at least the VH and/or VL regions and correlating said Complementarity Determining Regions.

25 12. A method according to claim 11, said Complementarity Determining Regions being selected from the group consisting of CDR1, CDR2 and CDR3.

13. A method according to any of claims 4-12, said correlation step (ii) additionally correlating at least one of the group consisting of: the strain of *Clostridium difficile*

- 49 -

infecting said at least one patient, the time point at which said B cells are isolated during infection of said at least one patient by *Clostridium difficile*, the age of said at least one patient, the sex of said at least one patient, and the race of said at least one patient.

5 14. A method according to any of claims 4-13, said B cells having been isolated from said at least one patient at a plurality of time points during infection of said at least one patient by *Clostridium difficile*, said correlation step (ii) correlating the time point during infection of said at least one patient by *Clostridium difficile* at which said B cells are isolated.

10 15. A method according to any of claims 4-13, said B cells having been isolated from at least two patients, at least one of whom has recovered from infection by *Clostridium difficile*, and at least one of whom has not recovered from infection by *Clostridium difficile*, said correlation step (ii) correlating the recovery of said at least two patients from
15 infection by *Clostridium difficile*.

16. A method according to any of claims 4-13, said B cells having been isolated from at least two patients, said patients being infected by different strains of *Clostridium difficile* producing said at least one antigen, said correlation step (ii) correlating said sequenced at
20 least the VH and/or VL coding regions of said B cells to identify a set of candidate sequences for antibodies, each of which is specific against at least one shared antigen produced by said different strains of *Clostridium difficile* or is specific against different antigens produced by said different strains of *Clostridium difficile*.

25 17. A method of manufacture of a medicament for the treatment of an infection by *Clostridium difficile* which produces at least one antigen, comprising the steps of:
 (i) performing a method according to any of claims 4-16; and

- 50 -

(ii) synthesising at least one antibody comprising at least one candidate sequence specific against said at least one antigen produced by *Clostridium difficile*.

18. A method of manufacture of a medicament according to claim 17, comprising the step of combining said synthesised at least one antibody with a pharmaceutically acceptable carrier, diluent or excipient.

19. A method of treatment of an infection of a patient by *Clostridium difficile* which produces at least one antigen, comprising the steps of:

- 10 (i) performing a method according to any of claims 4-16;
- (ii) synthesising at least one antibody comprising at least one candidate sequence specific against said at least one antigen produced by *Clostridium difficile*; and
- 15 (iii) administering a therapeutically effective quantity of said at least one synthesised antibody to said patient.

20. A method of producing a database which identifies candidate sequences for antibodies specific against at least one antigen produced by *Clostridium difficile*, comprising the steps of:

- 20 (i) performing a method according to any of claims 4-16; and
- (ii) storing the data produced by said method in said database.

21. A method of generating a report which identifies candidate sequences for antibodies specific against at least one antigen produced by *Clostridium difficile*, comprising the steps of:

- 25 (i) performing a method according to any of claims 4-16; and
- (ii) producing a report comprising the data produced by said method.

- 51 -

22. A method for determining the efficacy of a vaccine, comprising the steps of:

(i) with B cells isolated from at least one patient who has been administered said vaccine, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells; and

(ii) correlating said sequenced at least the CDR3 region of the VH and/or VL coding regions of said B cells to identify a set of candidate sequences for at least the CDR3 region of antibodies specific against said vaccine, each of said set of CDR3 candidate sequences or a sequence having at least 80% homology therewith occurring in total at a frequency of at least 1 percent in the set of sequences determined at step (i).

23. A method according to claim 22, correlation step (ii) comprising correlating said sequenced at least a CDR3 region of the VH and/or VL coding regions of said B cells with sequenced at least a CDR3 region of the VH and/or VL coding regions of B cells isolated from at least one patient who has been infected with *Clostridium difficile* against which vaccination with said vaccine is intended to stimulate a protective immune response.

24. A method according to either of claims 22 or 23, said correlation step (ii) additionally correlating at least one of the group consisting of: the time since administration of said vaccine to said at least one patient, the age of said at least one patient, the sex of said at least one patient, the race of said at least one patient, and the Complementarity Determining Regions of said sequenced at least the VH and/or VL coding regions.

25. A method according to any of claims 22-24, said method determining the efficacy of said vaccine in stimulating a protective immune response against *Clostridium difficile* against which vaccination with said vaccine is intended to stimulate a protective immune response.

- 52 -

26. A method of producing a database which identifies the efficacy of a vaccine, comprising the steps of:

- (i) performing a method according to any one of claims 22-25; and
- (ii) storing the data produced by said method in said database.

5

27. A method of generating a report which identifies the efficacy of a vaccine, comprising the steps of:

- (i) performing a method according to any one of claims 22-25; and
- (ii) producing a report comprising the data produced by said method.

10

28. A diagnostic test method for identifying a *Clostridium difficile* infection in a patient, comprising the steps of:

- (i) with B cells isolated from said patient, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells;
- 15 (ii) comparing said sequenced at least said CDR3 region of the VH and/or VL coding regions of said B cells with a set of sequences for at least the CDR3 region of antibodies specific against *Clostridium difficile*, and determining whether each of said set of CDR3 sequences or a sequence having at least 80% homology therewith occurs in total at a frequency of at least 1 percent
- 20 in the set of sequences determined at step (i); and
- (iii) correlating the results of comparison step (ii) to determine the presence or absence of a *Clostridium difficile* infection in said patient.

20

29. A diagnostic test method for determining the susceptibility of a patient to *Clostridium difficile* infection, comprising the steps of:

25

- (i) with B cells isolated from said patient, sequencing at least the CDR3 region of the VH and/or VL coding regions of said B cells;

- 53 -

- (ii) comparing said sequenced at least said CDR3 region of the VH and/or VL coding regions of said B cells with a set of sequences for at least the CDR3 region of antibodies specific against *Clostridium difficile*, and determining whether each of said set of CDR3 sequences or a sequence having at least 80% homology therewith occurs in total at a frequency of at least 1 percent in the set of sequences determined at step (i); and
- (iii) correlating the results of comparison step (ii) to determine the susceptibility of said patient to *Clostridium difficile* infection.

30. A diagnostic test kit for performing a diagnostic test method according to any one of claims 28-29.

- 1 -

SEQUENCE LISTING

<110> NeuTec Pharma PLC

5 <120> Clostridium difficile Focussed Antibodies

<130> GDM/JR-MP100462-WO

<150> GB0309126.1

10 <151> 2003-04-17

<160> 97

<170> PatentIn version 3.2

15 <210> 1

<211> 24

<212> DNA

<213> Artificial Sequence

20 <220>

<223> PCR Primer

<400> 1

25 gtccaccttg gtgttgctgg gctt 24

<210> 2

<211> 24

30 <212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

35 <400> 2

agactctccc ctgttgaagc tctt 24

- 2 -

<210> 3
<211> 24
<212> DNA
5 <213> Artificial Sequence

<220>
<223> PCR Primer

10 <400> 3
tgaggagacg gtgaccaggg tgcc 24

<210> 4
15 <211> 24
<212> DNA
<213> Artificial Sequence

<220>
20 <223> PCR Primer

<400> 4
tgaagagacg gtgaccattg tccc 24

25

<210> 5
<211> 24
<212> DNA
<213> Artificial Sequence

30

<220>
<223> PCR Primer

<400> 5
35 tgaggagacg gtgaccaggg ttcc 24

- 3 -

<210> 6
<211> 24
<212> DNA
<213> Artificial Sequence
5
<220>
<223> PCR Primer

<400> 6
10 tgaggagacg gtgaccgtgg tccc 24

<210> 7
<211> 23
15 <212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer
20

<400> 7
caggtgcagc tggcgcagtc tgg 23

25 <210> 8
<211> 23
<212> DNA
<213> Artificial Sequence

30 <220>
<223> PCR Primer

<400> 8
35 caggtgaact taagggagtc tgg 23

<210> 9

- 4 -

<211> 23
<212> DNA
<213> Artificial Sequence

5 <220>
<223> PCR Primer

<400> 9
caggtgcagc tggaggagtc tgg 23

10

<210> 10
<211> 23
<212> DNA
15 <213> Artificial Sequence

<220>
<223> PCR Primer

20 <400> 10
caggtgcagc tgcaggagtc ggg 23

<210> 11
25 <211> 23
<212> DNA
<213> Artificial Sequence

<220>
30 <223> PCR Primer

<400> 11
caggtgcagc tacagcagtg ggg 23

35

<210> 12
<211> 23

- 5 -

<212> DNA
<213> Artificial Sequence

<220>
5 <223> PCR Primer

<400> 12
gaggtgcagc tgttcagtc tgc 23

10

<210> 13
<211> 23
<212> DNA
<213> Artificial Sequence

15

<220>
<223> PCR Primer

<400> 13
20 caggtacagc tgcagcagtc agg 23

<210> 14
<211> 24
25 <212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

30

<400> 14
acgtttgatt tccaccttgg tccc 24

35

<210> 15
<211> 24
<212> DNA

- 6 -

<213> Artificial Sequence

<220>

<223> PCR Primer

5 <400> 15
acgtttgatc tccagcttgg tccc 24

10 <210> 16
<211> 24
<212> DNA
<213> Artificial Sequence

15 <220>
<223> PCR Primer

<400> 16
acgtttgata tccactttgg tccc 24

20

<210> 17
<211> 24
<212> DNA

25 <213> Artificial Sequence

<220>
<223> PCR Primer

30 <400> 17
acgtttgatc tccaccttgg tccc 24

<210> 18

35 <211> 24
<212> DNA
<213> Artificial Sequence

- 7 -

<220>

<223> PCR Primer

<400> 18

5 acgtttatttc tccagtcgtg tccc 24

<210> 19

<211> 23

10 <212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

15 <400> 19
gacatccaga tgacccagtc tcc 23

20 <210> 20

<211> 23

<212> DNA

<213> Artificial Sequence

25 <220>

<223> PCR Primer

<400> 20

30 gatgttgtga tgactcagtc tcc 23

<210> 21

<211> 23

<212> DNA

35 <213> Artificial Sequence

<220>

- 8 -

<223> PCR Primer

<400> 21
gaaattgtgt tgacgcagtc tcc 23

5

<210> 22
<211> 23
<212> DNA
10 <213> Artificial Sequence

<220>
<223> PCR Primer

15 <400> 22
gacatcgtga tgacccagtc tcc 23

<210> 23
20 <211> 23
<212> DNA
<213> Artificial Sequence

<220>
25 <223> PCR Primer

<400> 23
gaaactacac tcacgcagtc tcc 23

30

<210> 24
<211> 23
<212> DNA
<213> Artificial Sequence

35

<220>
<223> PCR primer

- 9 -

<400> 24
gaaattgtgc tgactcagtc tcc 23

5 <210> 25
<211> 17
<212> DNA
<213> Artificial Sequence

10 <220>
<223> PCR Primer

<400> 25
gtaaaacgac ggccagt 17

15

<210> 26
<211> 16
<212> DNA
20 <213> Artificial Sequence

<220>
<223> PCR Primer

25 <400> 26
aacagctatg accatg 16

<210> 27
30 <211> 16
<212> PRT
<213> Homo sapiens

<400> 27

35

Glu Ile Arg Ala Pro Asp His His Asp Phe Ser Gly Tyr Leu Gly Arg
1 5 10 15

- 10 -

<210> 28

<211> 9

<212> PRT

5 <213> Homo sapiens

<400> 28

Asp Ile Ser Ala Gly Gly Leu Asp Val

10 1 5

<210> 29

<211> 16

15 <212> PRT

<213> Homo sapiens

<400> 29

20 Asn Val Gly Ser Gly Ser Tyr Tyr Thr Gly Gly His Trp Phe Asp Pro

1 5 10 15

<210> 30

25 <211> 18

<212> PRT

<213> Homo sapiens

<400> 30

30

Asp Gly Val Arg Gln Tyr Ser Gly Gly Arg Tyr Ser Asn His Gly Met

1 5 10 15

35 Asp Val

- 11 -

<210> 31
<211> 10
<212> PRT
<213> Homo sapiens

5

<400> 31

Leu Thr Ala Ala Gly Gly His Phe Asp Pro
1 5 10

10

<210> 32
<211> 10
<212> PRT
<213> Homo sapiens

15

<400> 32

Asn Ser Arg Asp Ser Thr Gly Asn Gln Leu
1 5 10

20

<210> 33
<211> 11
<212> PRT
<213> Homo sapiens

25

<400> 33

Ala Ala Trp Asp Asp Ser Leu Ser Glu Phe Leu
1 5 10

30

<210> 34
<211> 9
<212> PRT
<213> Homo sapiens

35

- 12 -

<400> 34

Gln Gln Tyr Lys Gly Tyr Pro Leu Thr
1 5

5

<210> 35

<211> 16

<212> PRT

10 <213> Homo sapiens

<400> 35

Glu Ile Arg Ala Pro Asp His His Asp Leu Ser Gly Tyr Leu Gly Arg
15 1 5 10 15

<210> 36

<211> 16

20 <212> PRT

<213> Homo sapiens

<400> 36

25 Glu Ile Arg Ala Pro Asn His His Asp Phe Ser Gly Tyr Leu Gly Arg
1 5 10 15

<210> 37

30 <211> 16

<212> PRT

<213> Homo sapiens

<400> 37

35

Glu Val Arg Ala Pro Asp His His Asp Phe Ser Gly Tyr Leu Gly Arg
1 5 10 15

- 13 -

<210> 38
<211> 16
<212> PRT
5 <213> Homo sapiens

<400> 38

Glu Ile Arg Ala Pro Asp His His Asp Phe Ser Gly Tyr Leu Gly Cys
10 1 5 10 15

<210> 39
<211> 9
15 <212> PRT
<213> Homo sapiens

<400> 39

20 Asp Val Ser Ala Gly Gly Leu Asp Val
1 5

<210> 40
25 <211> 9
<212> PRT
<213> Homo sapiens

<400> 40
30

Glu Ile Ser Ala Gly Ala Leu Asp Val
1 5

35 <210> 41
<211> 16
<212> PRT

- 14 -

<213> Homo sapiens

<400> 41

5 Asp Val Gly Ser Gly Ser Tyr Tyr Thr Gly Gly His Trp Phe Asp Pro
1 5 10 15

<210> 42

10 <211> 16

<212> PRT

<213> Homo sapiens

<400> 42

15

Asn Val Gly Ser Gly Ser Tyr Tyr Thr Gly Gly His Trp Phe Glu Pro
1 5 10 15

20 <210> 43

<211> 16

<212> PRT

<213> Homo sapiens

25 <400> 43

Asn Val Gly Ser Gly Ser Tyr Tyr Thr Gly Gly His Trp Leu Asp Pro
1 5 10 15

30

<210> 44

<211> 16

<212> PRT

<213> Homo sapiens

35

<400> 44

- 15 -

Asn Val Gly Ser Gly Ser Tyr Tyr Thr Gly Gly His Trp Phe Asp Thr
1 5 10 15

5 <210> 45
<211> 16
<212> PRT
<213> Homo sapiens

10 <400> 45

Asn Val Gly Ser Gly Ser Tyr Tyr Thr Gly Gly His Trp Phe Gly Pro
1 5 10 15

15 <210> 46
<211> 15
<212> PRT
<213> Homo sapiens

20 <400> 46

Asn Val Gly Ser Gly Ser Tyr Tyr Thr Ala Thr Cys Phe Asp Pro
1 5 10 15

25 <210> 47
<211> 17
<212> PRT
30 <213> Homo sapiens

<400> 47

Gly Ala Lys His Phe Glu Gly Ser Gly Ser Trp Phe Ser Trp Phe Asp
35 1 5 10 15

- 16 -

Pro

5 <210> 48
 <211> 11
 <212> PRT
 <213> Homo sapiens

10 <400> 48

Asn Ser Arg Asp Asn Thr Gly His His Val Val
1 5 10

15 <210> 49
 <211> 11
 <212> PRT
 <213> Homo sapiens

20 <400> 49

Asn Ser Arg Asp Ser Ser Gly Asn His Val Val
1 5 10

25 <210> 50
 <211> 11
 <212> PRT
30 <213> Homo sapiens

<400> 50

Ser Ser Arg Asp Asn Ser Gly Asp Arg Tyr Val
35 1 5 10

- 17 -

<210> 51
<211> 10
<212> PRT
<213> Homo sapiens

5

<400> 51

Asn Ser Arg Asp Gly Thr Gly Asn Gln Leu
1 5 10

10

<210> 52
<211> 10
<212> PRT
15 <213> Homo sapiens

<400> 52

Asn Ser Arg Asp Thr Asn Gly Asp Gln Leu
20 1 5 10

<210> 53
<211> 11
25 <212> PRT
<213> Homo sapiens

<400> 53

30 Asn Ser Arg Asp Ser Ser Gly Tyr His Val Ile
1 5 10

<210> 54
35 <211> 11
<212> PRT
<213> Homo sapiens

- 18 -

<400> 54

Ser Ser Arg Asp Ser Lys Gly His Arg Tyr Val
1 5 10

5

<210> 55

<211> 11

<212> PRT

10 <213> Homo sapiens

<400> 55

Ser Ser Arg Asp Ser Asn Gly Asn Arg Tyr Val
1 5 10

15

<210> 56

<211> 11

20 <212> PRT

<213> Homo sapiens

<400> 56

Ser Ser Arg Asp Thr Lys Gly His Arg Tyr Val
1 5 10

25

<210> 57

30 <211> 10

<212> PRT

<213> Homo sapiens

<400> 57

35

Lys Pro Arg Asp Ser Ser Gly Asn His Val
1 5 10

- 19 -

<210> 58
<211> 11
<212> PRT
5 <213> Homo sapiens

<400> 58

Thr Ser Arg Asp Ser Asn Gly Asn Arg Tyr Val
10 1 5 10

<210> 59
<211> 11
15 <212> PRT
<213> Homo sapiens

<400> 59

Ala Ala Trp Asp Asp Asn Leu Ser Ala Tyr Val
20 1 5 10

<210> 60
25 <211> 12
<212> PRT
<213> Homo sapiens

<400> 60

30 Ala Ala Trp Asp Asp Ser Leu Gly Gly Lys Tyr Val
1 5 10

35 <210> 61
<211> 11
<212> PRT

- 20 -

<213> Homo sapiens

<400> 61

5 Ala Ala Trp Asp Asp Ser Leu Asn Gly Val Val
1 5 10

<210> 62

10 <211> 11

<212> PRT

<213> Homo sapiens

<400> 62

15 Ala Ala Trp Asp Asp Ser Val Lys Gly Trp Val
1 5 10

20 <210> 63

<211> 11

<212> PRT

<213> Homo sapiens

25 <400> 63

Ala Ala Trp Asp Asp Ser Val Arg Gly Trp Val
1 5 10

30

<210> 64

<211> 11

<212> PRT

<213> Homo sapiens

35

<400> 64

- 21 -

Ala Ala Trp Asp Asp Ser Val Arg Ser Trp Val
1 5 10

5 <210> 65
<211> 11
<212> PRT
<213> Homo sapiens

10 <400> 65

Ala Ser Trp Asp Asp Ser Leu Asn Gly Val Val
1 5 10

15 <210> 66
<211> 11
<212> PRT
<213> Homo sapiens

20 <400> 66

Ala Ser Trp Asp Asp Ser Gln Ala Ala Leu Val
1 5 10

25 <210> 67
<211> 11
<212> PRT
30 <213> Homo sapiens

<400> 67

Ser Ala Trp Asp Ser Ser Leu Ser Thr Trp Val
35 1 5 10

- 22 -

<210> 68
<211> 11
<212> PRT
<213> Homo sapiens

5

<400> 68

Gln Ser Tyr Asp Ser Ser Leu Ser Gly Trp Val
1 5 10

10

<210> 69
<211> 10
<212> PRT
<213> Homo sapiens

15

<400> 69

Ser Ser Tyr Thr Asn Thr Asn Pro Tyr Val
1 5 10

20

<210> 70
<211> 9
<212> PRT
<213> Homo sapiens

25

<400> 70

Gln Gln Phe Lys Ser Phe Pro Leu Thr
1 5

30

<210> 71
<211> 9
<212> PRT
<213> Homo sapiens

35

- 23 -

<400> 71

Gln Gln Phe Gln Ser Tyr Pro Val Thr
1 5

5

<210> 72

<211> 9

<212> PRT

10 <213> Homo sapiens

<400> 72

Gln Gln Tyr Lys Ser Tyr Pro Leu Thr
1 5

15

<210> 73

<211> 9

20 <212> PRT

<213> Homo sapiens

<400> 73

25 His Gln Tyr Tyr Arg Phe Pro Leu Thr
1 5

<210> 74

30 <211> 9

<212> PRT

<213> Homo sapiens

<400> 74

35

Leu Gln Tyr Lys Lys Trp Pro Leu Thr
1 5

- 24 -

<210> 75
<211> 9
<212> PRT
5 <213> Homo sapiens

<400> 75

Gln His Phe Ser His Tyr Pro Leu Thr
10 1 5

<210> 76
<211> 9
15 <212> PRT
<213> Homo sapiens

<400> 76

Gln His Tyr Tyr Ser Tyr Pro Leu Thr
20 1 5

<210> 77
25 <211> 18
<212> PRT
<213> Homo sapiens

<400> 77
30
Asp Gly Val Arg Gln Tyr Asn Gly Gly Arg Tyr Ser Asn His Gly Met
1 5 10 15

35 Asp Val

- 25 -

<210> 78
<211> 18
<212> PRT
<213> Homo sapiens

5

<400> 78

Asp Gly Val Arg Gln Tyr Ser Gly Gly Lys Tyr Ser Asn His Gly Met
1 5 10 15

10

Asp Val

15

<210> 79
<211> 18
<212> PRT
<213> Homo sapiens

20

<400> 79

Asp Gly Val Arg Gln Tyr Ser Gly Gly Arg Tyr Ser Asn His Gly Met
1 5 10 15

25

Asn Val

30

<210> 80
<211> 18
<212> PRT
<213> Homo sapiens

35

<400> 80

- 26 -

Asp Gly Val Ser Gln His Asn Gly Gly Arg Tyr Ser Asn His Gly Met
1 5 10 15

5 Asp Val

10 <210> 81
<211> 18
<212> PRT
<213> Homo sapiens

15 <400> 81

Asp Gly Val Ser Gln Tyr Ser Gly Gly Arg Tyr Ser Asn His Gly Met
1 5 10 15

20 Asp Val

25 <210> 82
<211> 18
<212> PRT
<213> Homo sapiens

30 <400> 82

Asn Gly Val Arg Gln Tyr Ser Gly Gly Arg Tyr Ser Asn His Arg Met
1 5 10 15

35 Asp Val

- 27 -

<210> 83
<211> 18
<212> PRT
<213> Homo sapiens

5

<400> 83

Glu Glu Glu Arg Gln Tyr Ser Gly Gly Arg Tyr Ser Asn His Gly Met
1 5 10 15

10

Asp Val

15

<210> 84
<211> 18
<212> PRT
<213> Homo sapiens

20

<400> 84

Asp Gly Val Arg Gln Tyr Ser Gly Ala Asp Thr Pro Asn His Gly Met
1 5 10 15

25

Asp Val

30

<210> 85
<211> 10
<212> PRT
<213> Homo sapiens

35

<400> 85

- 28 -

Leu Thr Ala Ala Gly Ala His Phe Asp Pro
1 5 10

5 <210> 86
<211> 10
<212> PRT
<213> Homo sapiens

10 <400> 86

Leu Thr Ala Ala Gly Gly Arg Phe Asp Pro
1 5 10

15 <210> 87
<211> 11
<212> PRT
<213> Homo sapiens

20 <400> 87

His Ser Arg Asp Asp Thr His Tyr Pro Val Ile
1 5 10

25 <210> 88
<211> 10
<212> PRT
30 <213> Homo sapiens

<400> 88

Asn Ser Arg Asp Ser Asn Asn His Val Val
35 1 5 10

- 29 -

<210> 89
<211> 9
<212> PRT
<213> Homo sapiens

5

<400> 89

Gln Ser Arg Asn Asn Thr Tyr Val Leu
1 5

10

<210> 90
<211> 11
<212> PRT
15 <213> Homo sapiens

<400> 90

His Thr Arg Asp Thr Leu Phe Pro Val Asp Phe
20 1 5 10

<210> 91
<211> 19
25 <212> PRT
<213> Homo sapiens

<400> 91

30 Asp Met Gly Met Phe Cys Ser Gly Val Ile Cys Tyr Asp Tyr Tyr Gly
1 5 10 15

Met Asp Val

35

- 30 -

<210> 92
<211> 4
<212> PRT
<213> Artificial Sequence

5

<220>
<223> Used for defining CDR region

10 <220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Any amino acid

15 <400> 92

Phe Gly Xaa Gly
1

20

<210> 93
<211> 4
<212> PRT
<213> Artificial Sequence

25

<220>
<223> Used for defining CDR region

30 <220>
<221> MISC_FEATURE
<222> (2)..(4)
<223> Any amino acid

35 <400> 93

Cys Xaa Xaa Xaa

- 31 -

1

<210> 94

5 <211> 5

<212> PRT

<213> Artificial Sequence

<220>

10 <223> Used for defining CDR region

<400> 94

Leu Glu Trp Ile Gly

15 1 5

<210> 95

<211> 4

20 <212> PRT

<213> Artificial Sequence

<220>

<223> Used for defining CDR region

25

<220>

<221> MISC_FEATURE

<222> (3)..(3)

30 <223> Any amino acid

<400> 95

Trp Gly Xaa Gly

35 1

- 32 -

<210> 96
<211> 10
<212> PRT
<213> Homo sapiens

5

<400> 96

Gln Ser Tyr Asp Ser Ser Arg Gly Arg Val
1 5 10

10

<210> 97
<211> 9
<212> PRT
<213> Homo sapiens

15

<400> 97

Met Gln Gly Ile Arg Pro Pro Arg Thr
1 5

20

INTERNATIONAL SEARCH REPORT

In national Application No

T/GB2004/001619

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K16/12 A61K39/40 C12Q1/68 G01N33/563

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBL, EMBASE, MEDLINE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 197 39 685 A (EICHEL STREIBER CHRISTOPH VON) 11 March 1999 (1999-03-11) page 3, line 1 - line 11 examples 1-6	1-3, 30
X	WO 00/12562 A (GENENTECH INC) 9 March 2000 (2000-03-09) figure 2 claim 8 SEQ ID No. 15	1-3

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

2 September 2004

Date of mailing of the international search report

27/09/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Ulbrecht, M

INTERNATIONAL SEARCH REPORT

International Application No

T/GB2004/001619

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97/20932 A (ALLEN DEBORAH JULIE ; CAMBRIDGE ANTIBODY TECH (GB); MCCAFFERTY JOHN GE) 12 June 1997 (1997-06-12) example 1 table 1 figure 1b claim 14	1-3, 30
X	----- WHITE HARRY ET AL: "Analysis of immunoglobulin (Ig) isotype diversity and IgM/D memory in the response to phenyl-oxazolone" JOURNAL OF EXPERIMENTAL MEDICINE, vol. 191, no. 12, 19 June 2000 (2000-06-19), pages 2209-2219, XP002293496 ISSN: 0022-1007 page 2210, left-hand column, paragraph 3 - page 2211, left-hand column, paragraph 1 -----	30
X	NIE XIAOBO ET AL: "Immunization with immune complex alters the repertoire of antigen-reactive B cells in the germinal centers" EUROPEAN JOURNAL OF IMMUNOLOGY, vol. 27, no. 12, December 1997 (1997-12), pages 3517-3525, XP009035552 ISSN: 0014-2980 page 3524, left-hand column, paragraph 3 - right-hand column, paragraph 1 -----	30
P, X	WO 03/052416 A (MATTHEWS RUTH CHRISTINE ; NEUTEC PHARMA PLC (GB); RIGG GORDON PATRICK) 26 June 2003 (2003-06-26) the whole document in particular SEQ ID Nos. 37 and 50.	1, 3-27, 30
P, X	----- HOLT L J ET AL: "Domain antibodies: proteins for therapy" TRENDS IN BIOTECHNOLOGY, ELSEVIER PUBLICATIONS, CAMBRIDGE, GB, vol. 21, no. 11, November 2003 (2003-11), pages 484-490, XP004467495 ISSN: 0167-7799 the whole document -----	1-3, 17-19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB2004/001619

Box No. 1 Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:

a. type of material

☒

a sequence listing

☐

table(s) related to the sequence listing

b. format of material

☒

in written format

☒

in computer readable form

c. time of filing/furnishing

☒

contained in the international application as filed

☐

filed together with the international application in computer readable form

☐

furnished subsequently to this Authority for the purpose of search

2. ☐ In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

INTERNATIONAL SEARCH REPORT

international application No.
PCT/GB2004/001619

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 19
because they relate to subject matter not required to be searched by this Authority, namely:
Although claim 19 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

T/GB2004/001619

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 19739685	A	11-03-1999	DE 19739685 A1	11-03-1999
			AT 254139 T	15-11-2003
			AU 9742698 A	29-03-1999
			BR 9815367 A	06-11-2001
			CA 2303202 A1	18-03-1999
			CN 1273588 T	15-11-2000
			DE 59810172 D1	18-12-2003
			WO 9912971 A2	18-03-1999
			EP 0994904 A2	26-04-2000
			ES 2210832 T3	01-07-2004
			JP 2001515920 T	25-09-2001
			US 6667035 B1	23-12-2003
			US 2004137601 A1	15-07-2004
WO 0012562	A	09-03-2000	AT 217889 T	15-06-2002
			AU 766551 B2	16-10-2003
			AU 5692399 A	21-03-2000
			BR 9913435 A	25-09-2001
			CA 2341276 A1	09-03-2000
			CN 1317015 T	10-10-2001
			DE 69901569 D1	27-06-2002
			DE 69901569 T2	19-12-2002
			DK 1107996 T3	16-09-2002
			EP 1107996 A1	20-06-2001
			ES 2177316 T3	01-12-2002
			JP 2002523081 T	30-07-2002
			NZ 509430 A	26-04-2002
			PT 1107996 T	31-10-2002
			WO 0012562 A1	09-03-2000
			US 6624295 B1	23-09-2003
			ZA 200100681 A	24-01-2002
WO 9720932	A	12-06-1997	AT 200516 T	15-04-2001
			AU 703319 B2	25-03-1999
			AU 1103697 A	27-06-1997
			CA 2239519 A1	12-06-1997
			DE 69612509 D1	17-05-2001
			DE 69612509 T2	18-10-2001
			DK 865492 T3	11-06-2001
			EP 0865492 A1	23-09-1998
			ES 2157473 T3	16-08-2001
			WO 9720932 A1	12-06-1997
			JP 2000504204 T	11-04-2000
			US 5872215 A	16-02-1999
WO 03052416	A	26-06-2003	CA 2471570 A1	26-06-2003
			EP 1415002 A2	06-05-2004
			WO 03052416 A2	26-06-2003